

“STUDY OF LIPID PROFILE IN PRE-ECLAMPSIA.”

¹PRADNYA PHALAK*, MONA TILAK*,

*Departments of Biochemistry,

Pad. Dr. D. Y. Patil Medical College, D. Y. Patil University, Pimpri, Pune.

Corresponding author:Dr. Pradnya Phalak; Email ID: pradnyaphalak@gmail.com

ABSTRACT:

Introduction: Pre-eclampsia is common medical complication of pregnancy. The association of altered lipid profile in essential hypertension is well documented. The present study was planned to study the role of altered lipid profile in the development of pre-eclampsia.

Method: The study comprised of 30 normal pregnant women and 30 pre-eclamptic women in their third trimester of pregnancy. Serum Total cholesterol, Triglycerides, LDL Cholesterol and HDL- Cholesterol by enzymatic colorimetric method were done.

Results: There was significant rise in Serum Total cholesterol, Triglycerides, LDL Cholesterol and a significant decrease in HDL Cholesterol in pre-eclamptic group as compared to normal healthy pregnant women.

Conclusion: Altered lipid profile also has a potential role in the genesis of endothelial dysfunction and expression of pre-eclampsia. Early detection of these parameters may help patient by preventing complications in pre-eclampsia and is going to aid in better management of Pre-eclampsia.

Key words: Lipid Profile, Dyslipidemia, Pre-eclampsia.

INTRODUCTION:

Pre-eclampsia is common medical complication of pregnancy. In India, the incidence of pre-eclampsia is reported to be 8-10% of the pregnancies[1]. It contributes significantly to maternal and fetal mortality and morbidity. Pre-eclampsia is a multisystem disorder characterized by hypertension to the extent of 140/90 mm Hg or more, proteinuria (≥ 300 mg/day) and edema induced by pregnancy after 20th week[2]. Without intervention, pre-eclampsia progresses to eclampsia which is characterized by malignant hypertension and epileptiform convulsions requiring emergency caesarian section[3].

The association of altered lipid profile in essential hypertension is well documented. In early pregnancy, there is increased body fat accumulation associated with increased lipogenesis, while in late pregnancy, there is accelerated breakdown of fat depots which play an

important role in fetal development (4). Early pregnancy dyslipidemia is associated with an increased risk of pre-eclampsia(5).

Women with a history of pre-eclampsia have significant differences in lipid parameters and an increased susceptibility to lipoprotein oxidation when compared with women who had normal pregnancy(6). Therefore, simple measurement of serum lipid profile may be of good predictive value in pre-eclampsia. With this background in mind the present study was conducted to assess lipid profile in cases of pre-eclampsia and normal healthy pregnant women.

MATERIALS AND METHODS: The present study was carried out in the Department of Biochemistry in collaboration with Department of Obstetrics and Gynecology, B. J. Medical College and Sasson General Hospitals, Pune. Informed consent was taken from all individual subjects inducted into the study. The study comprised of 30 normal

healthy pregnant women and 30 pre-eclampsia cases attending antenatal OPD in their third trimester of pregnancy.

The diagnosis of pre-eclampsia was based on the definition of American College of Obstetrics and Gynecologists (7).1) Systolic blood pressure greater than 140 mm Hg or a rise of at least 30 mmHg or 2) Diastolic blood pressure greater than 90 mm Hg or a rise of at least 15 mmHg (manifested on two occasions at least 6 hours apart) and 3) Proteinuria of 300 mg or greater in 24 hours urine collection or protein concentration of 1 gm/L (on two occasions of at least 6 hours apart).

Inclusion criteria for controls: Primigravida with normal BP, no proteinuria and without any other systemic or endocrine disorder. They were age matched with the cases. All subjects included were in their third trimester (gestational age of ≥ 24 weeks).

Exclusion criteria included diabetes mellitus with or without treatment, obesity, severe anemia (Hb < 6 gm%) or subjects suffering from any other systemic or endocrine disorder. Fasting blood samples (8ml) were collected by venipuncture and analysed for Serum Triglycerides (TG), Total cholesterol (TC), HDL cholesterol (HDL-C), LDL cholesterol (LDL-C) by the following methods.

Estimation of Serum Cholesterol: Serum total cholesterol was estimated by cholesterol oxidase/ Peroxidase (CHOD/POD) colorimetric endpoint method, using Liquizyme kit.

Cholesterol esters are hydrolysed by cholesterol esterase to produce free cholesterol and fatty acid. This free cholesterol gets oxidized in the presence of cholesterol oxidase to liberate cholestenone and hydrogen peroxide. Liberated hydrogen peroxide combines with hydroxyl benzoate and 4

aminoantipyrine in the presence of peroxidase to form red coloured quinonimine complex, the intensity of which is measured at 505nm (490- 530nm).

Estimation of Serum HDL-C: Serum HDL-C was estimated by Direct method using, Liquizyme kit (NCCLS, 1999). The system utilizes a combination of surfactants, phosphoric acid, organic acids, and inorganic acids, specifically binding LDL-C, VLDL-C and chylomicrons. Only HDL cholesterol is detected by the enzymatic CHOD/ POD method.

Estimation of Serum Triglycerides: Serum triglyceride was estimated by Glycerol phosphate oxidase/ Peroxidase (GPO/POD) colorimetric endpoint method using Liquizyme kit. Triglyceride is hydrolysed by lipoprotein lipase to free fatty acids and glycerol. Glycerol kinase converts glycerol to glycerol phosphate, which gets oxidized to dihydroxy acetone phosphate and hydrogen peroxide by glycerol phosphate oxidase. Hydrogen peroxide so formed reacts with 4 aminoantipyrine in the presence of peroxidase to give a purple coloured complex which is read at 546 nm.

Calculation of LDL-C and VLDL-C : VLDL-C level in serum is derived by dividing serum triglycerides by 5 and LDL-C is obtained using Friedwald's formula (Friedwald *et al.*, 1972), namely, $LDL-C = Total\ cholesterol - (HDL-C + VLDL-C)$.

Data were statistically analyzed by unpaired T test and expressed in terms of 'P' value.

RESULTS :

There was a significant rise in Serum Triglycerides ($149.75 \pm 20.42^{**}$, 125.20 ± 18.30) Total cholesterol ($195.40 \pm 15.11^{**}$, 176.34 ± 12.10) and LDL-C ($134.5 \pm 11.65^{**}$, 114.2 ± 11.32) levels and a significant decrease in HDL-C ($42.26 \pm 12.03^{**}$, 55.20 ± 10.02) levels in cases as compared to controls ($P < 0.0001$) (Table I).

Table I: Comparison of serum lipid profile in the study groups:

PARAMETER	STUDY GROUP	SAMPLE SIZE (N)	MEAN±SD (mg/dl)
Total Cholesterol	Patients	30	195.40±15.11**
	Controls	30	176.34±12.10
LDL	Patients	30	134.5±11.65**
	Controls	30	114.2±11.32
HDL	Patients	30	42.26±12.03**
	Controls	30	55.20±10.02
TG	Patients	30	149.75±20.42**
	Controls	30	125.20±18.30

**p<0.001-highly significant

DISCUSSION: Pre-eclampsia has long known to be associated with abnormal placentation and impaired placental perfusion. However, other conditions characterized by poor placentation, such as intrauterine growth retardation, do not necessarily result in pre-eclampsia(8). This has led to the growing concept that maternal predisposing factors must combine with the placental disorder to result in preeclamptic maternal syndrome (9). In the recent last years, several lines of evidence have suggested that part of this maternal predisposition could be explained by abnormal lipid metabolism. In this study, we observed significantly increased serum TC, TG and LDL-C and a decreased HDL-C in cases as compared to controls. Normal human pregnancy results in physiologic hyperlipidemia involving a gestational rise in blood triglyceride (TG) and cholesterol. There is a marked rise in

serum TG in normal pregnancy as compared to nonpregnant women, which may be as high as two to three folds in the third trimester (10).

The principle modulator of this hypertriglyceridemia is hyperoestrogenemia in pregnancy that induces hepatic biosynthesis of TGs (11). The anabolic phase of early pregnancy encourages lipogenesis and fat storage in preparation for rapid foetal growth in late pregnancy(12). Lipolysis is increased as a result of insulin resistance, leading to increased flux of fatty acids to the liver promoting the synthesis of very low-density lipoproteins and increased triglyceride (TG) concentrations(13).

During pregnancy, there is an increase in the hepatic lipase activity and decrease in lipoprotein lipase activity(14). Hepatic lipase is responsible for the increased synthesis of the TGs at the hepatic level, whereas the decreased activity of lipoprotein lipase is

responsible for the decreased catabolism at the adipose tissue level, the net effect of which will be an increase in circulating TGs (15).

Because of a decrease in the activity of lipoprotein lipase, very-LDL remains in the plasma for longer and leads to the accumulation of LDL. An increase in LDL is associated with the development of atherosclerosis(16). Women with preeclampsia display additional alterations in blood lipids reflecting disordered lipid and lipoprotein metabolism. Serum TG levels rose much more in pre- eclampsia as reported by other studies (5, 17) and as seen in our study also. Increased TG, found in pre-eclampsia is likely to be deposited in predisposed vessels, such as the uterine spiral arteries and contributes to the endothelial dysfunction (14).

There was a significant rise in TC levels in pre-eclampsia as compared to normal pregnancy in our study, which was similar to other reports (15, 16) however other studies reported no alteration in TC levels (8, 13). A significant rise in the LDL-C levels was seen in pre-eclampsia as compared to controls in our study and also by other workers (6,19). In our study there was significant fall in HDL-C in pre-eclampsia cases as compared to controls. Low HDL-C in pre-eclampsia is due to insulin resistance (17). According to Pirzado et al, there is a direct correlation between adipose tissue lipoprotein lipase activity and plasma HDL cholesterol(18). This direct correlation may be responsible for low levels of HDL cholesterol. Thus,

dyslipidemia mediated activation of the endothelial cells and the placentally derived endothelial disturbing factors like lipid peroxides could be a possible cause in the pathogenesis of PIH (20). This association may be significant in understanding the pathological process of pre-eclampsia and may help in developing strategies for prevention and early diagnosis of pre-eclampsia. Thus, estimation of lipid profile may have a predictive role in the assessment of the extent of endothelial damage in pre-eclampsia and may help patient by preventing or foreseeing the effects of complications in pre-eclampsia.

CONCLUSION: Hence early detection of these parameters is going to aid in better management of pre- eclampsia cases which is important to improve the maternal and fetal outcome in pre- eclampsia.

REFERENCES:

1. Kamath S. Hypertension in pregnancy. JAPL 2006; 54: 269-270
2. Dutta D.C (2001), Hypertensive disorders in pregnancy, in :Textbook of obstetrics, Ed. Konar, H.L., 5th edition,234-55
3. Packer CS. Biochemical markers and physiological parameters as indices for identifying patients at risk of developing pre-eclampsia. J Hypertens 2005;23:(1)45-6.
4. Herrera, E (2002). Lipid metabolism in pregnancy and its consequences in the fetus and newborn. Endocrine 19 43 – 55.
5. Enquobahrie DA, Williams MA, Butler CL, Frederick IO, Miller RS, Luthy DA. Maternal plasma lipid concentrations in early pregnancy and risk of preeclampsia. Am J Hypertens 2004; 17(7): 574–581.
6. Gratacos E, Casals E, Gomez O, Llorba E, Mercador I, Cararach V, Cabero L. Increased susceptibility to low density lipoprotein oxidation in women with a history of preeclampsia. Br J of Obstet Gynaecol 2003; 110(4): 400–4.
7. K. Yagi, Assay for lipid peroxide level and its clinical significance, in: Lipid Peroxides in Biology and Medicine, New York. 1982; 233–242.

8. Khong TY, DE Wolf, Robetson WB, Brosens I. Inadequate maternal vascular response to placentation in pregnancies complicated by pre-eclampsia and by small for gestational age infants. *Br J Obstet Gynaecol* 1986;93:1049-1059.
9. Broughton Pipkin F. What is the place of genetics in the pathogenesis of pre-eclampsia? *Biol Neonate* 1999;76:325-330.
10. Chiang,A.N.,Yang, M.L.,Hung, J.H., Chon, P., Shyn, S.K. and Ng,H.T.(1995).Alteration of serum lipid levels and their biological relevances during and after pregnancy.*Life sci.* 56(26),2367-75.
11. Glueck, C.J.,Fallet, R.W. and Scheel, D.(1975) Effects of oestrogenic compounds on triglyceride kinetics.*Metabolism* 24, 537-45.
12. Kaaja R. Insulin resistance syndrome in pre-eclampsia. *Sem Reprod Endocrinol* 1998; 16: 41–6.
13. Kaaja R., Tikkanen MJ. (1995) Serum lipoproteins, insulin and urinary prostanoid metabolites in normal and hypertensive pregnant women.*Obstet. Gynecol* 1995;85(3):253-6.
14. Patrizia B, Giancarlo T, Franca E, et al. Lipoprotein metabolism during normal pregnancy. *Am J Obstet Gynecol* 1999; 181(2): 430-4.
15. Rubina A, Tabassum M. Pre-eclampsia and lipid profile. *Pak J Med Sci* 2007; 23: 751–4.
16. Ross R. Atherosclerosis: an inflammatory disease. *N Engl J Med* 1999; 340: 115–26.
17. Cekmen, M.B.,Erbagci, A.B.(2003) plasma lipid and lipoprotein concentration in pregnancy induced hypertention,*Clin. Biochem.*36(7),575-8.
18. Pirzado ZA, Sangi SA, Malik R. High density lipoprotein cholesterol (HDL) metabolism and its role in the ischemic heart disease. *Pak J Med Res* 1999; 38: 38–41.
19. Wakatsuki A, Ikenoue N, Okatani Y, Shinohara K, Fukaya T. Lipoprotein particles in preeclampsia: Susceptibility to oxidative modification. *Obstet Gynecol* 2000; 96(1): 55–5
20. Lorensten, B. and Henriksen, T.(1998)Plasma lipids and vascular dysfunction in pre-eclampsia.

Source of support: Nil; Conflict of Interest: Nil
Date of submission: 2 Nov 2012
Date of Provisional Acceptance: 11 Nov 2012
Date of Peer review approval: 22 Nov 2012
Date of final drafting: 24 November 2012
Date of Publication: 2 December 2012