

**Original article:**

## **Adenosine Deaminase Levels as a Tool for the Diagnosis of Tuberculous Pleural Effusion: A Cross-Sectional Study**

**Ghubdee Ramakrishna Vishnupant**

Associate Professor, Department of Pulmonary Medicine, Smt. KashibaiNavale Medical College, Narhe, Pune, Maharashtra, India.

**Corresponding Author:** Dr. Ghubdee Ramakrishna Vishnupant, Associate Professor, Department of Pulmonary Medicine, Smt. KashibaiNavale Medical College, Narhe, Pune, Maharashtra, India.

---

### **Abstract**

**Introduction:** Tuberculous pleural effusion (TPE) is a common manifestation of extrapulmonary tuberculosis, and is the leading cause of pleural effusion in developing world regions, while it is much less common in developed countries. The aim of the study is to determine that the Adenosine Deaminase levels can use as a tool for the diagnosis of tubercular pleural effusion.

**Materials and Methods:** This cross-sectional study was conducted in one hundred ten cases of pleural effusion, in which seventy one cases were tubercular pleural effusion and thirty nine cases were non tubercular pleural effusion. A p-value < 0.05 was considered statistically significant. IBM SPSS Statistics 21 manufactured by IBM USA was used for entire calculations.

**Result:** In the present study the mean of ADA level was 173.3, LDH level was 184.91, T. Protein was 4.80, Glucose was 112.34 and the total cell count was 3996 in tuberculosis pleural effusion group. These variables are statistically significant (<0.05). This study has clearly shown that ADA levels are significantly high in patients with tubercular pleural effusion ( $173.3 \pm 34.93$  U/L) compared to that ( $20.46 \pm 7.34$  U/L) in non tuberculous group.

**Conclusion:** This study has clearly shown that ADA levels are significantly high in patients with tubercular pleural effusion compared to that in non tuberculous group.

**Keywords:** Tuberculous pleural effusion, Adenosine Deaminase, LDH.

---

### **Introduction**

Adenosine deaminase (ADA) is an enzyme which catalyses the conversion of adenosine to inosine and plays an important role in the differentiation of lymphoid cells. Activity is high in diseases in which cellular immunity is stimulated.<sup>1</sup> Different cut off values of ADA ranging from 30–100 IU/L have been used in various studies, with differing sensitivities and specificities.<sup>2</sup>

Tuberculous pleural effusion (TPE) is a common manifestation of extrapulmonary tuberculosis, and is the leading cause of pleural effusion in developing world regions, while it is much less common in

developed countries.<sup>3-5</sup> The diagnosis of pleural TB is still challenging in clinical practice. The diagnosis of TPE depends on the demonstration of tubercle bacilli in pleural fluid, a pleural biopsy specimen or sputum, or the demonstration of granulomas in the pleura.<sup>6</sup> Due to the paucity of Mycobacterium tuberculosis in the pleural fluid, the performance of a pleural biopsy has historically been considered the most reliable method to confirm the diagnosis when tuberculous aetiology of a pleural effusion is suspected. However, since pleural tissue sampling is more difficult than simple thoracentesis, pleural fluid markers of TPE have been extensively

evaluated as an attractive alternative to pleural biopsy.<sup>7</sup> ADA is the most cost effective pleural fluid marker and is routinely employed as a screening tool, in particular, in countries where tuberculosis is endemic.<sup>3</sup>

Recent estimates show that around 8-10 million new tuberculosis cases occur each year in the world and 2-3 million die.<sup>8</sup> In developing countries, TB is one of the common opportunistic infections in people who are seropositive for human immunodeficiency virus (HIV).<sup>9</sup> Tuberculosis is classified as pulmonary, extrapulmonary, or both.<sup>10</sup> Pleural tuberculosis accounts for fewer than 1% of all exudative effusions in Western countries, occurring in only 3% to 5% of tuberculosis patients and in developing countries like India, it is responsible for 30% to 80% of all pleural effusions encountered.<sup>11</sup>

Adenosine Deaminase Activity (ADA) in pleural fluid which identifies an inflammatory process triggered by MBT.<sup>12-14</sup> The most recent study, a meta-analysis by Lian QL and co-workers, shows a very good accuracy in measuring ADA in pleural effusion to diagnose TB pleural effusion. Nevertheless the best cutoff they found was 40 U/L given a positive likelihood (LH) of 9, 03 and negative LH ratio of 0,01 which are important values to consider.<sup>15</sup>

ADA is involved in the proliferation and differentiation of lymphocytes, especially T lymphocytes.<sup>16</sup> The increase in the ADA activity in patients with TB may indicate the cellular immune response and T lymphocyte activation in the disease.<sup>17</sup> T lymphocytes have ADA level 10 to 12 times higher than B lymphocytes. ADA activity varies depending on the proliferative status and maturity of cells.<sup>18</sup> The level of ADA is increased in TPE and this determination has acquired popularity

as a diagnostic test in the high incidence area of TPE, because ADA measurement is a less expensive, minimally invasive, rapid and readily accessible test.

The main aim of the study is to determine that the Adenosine Deaminase levels can be used as a tool for the diagnosis of tubercular pleural effusion.

#### **Materials and Methods**

This cross-sectional study was conducted in one hundred ten cases of pleural effusion, in which seventy one cases were tubercular pleural effusion and thirty nine cases were non tubercular pleural effusion. This study was done in the department of pulmonary medicine, Smt. Kashibai Navale Medical College, Narhe, Pune, Maharashtra, India. The study protocol was approved from the ethical committee of the college. Signed consent forms were obtained from all the subjects included in this study. Parameters studied were total cell count, differential cell count, glucose, total proteins, lactate dehydrogenase, and adenosine deaminase.

Thoracentesis was attempted at one inter space below the spot where tactile fremitus is lost and the percussion note becomes dull.<sup>19</sup> The exact location for the thoracentesis attempt was just superior to a rib.<sup>20,21</sup>

Pleural fluid was collected after the skin was cleaned with antiseptic solution. The sterile drape with the central hole was then taped to the patient's back and another sterile drape was placed on the bed. Anaesthetization of the skin, the periosteum of the rib and the parietal pleura was done.<sup>22</sup> 20/50 ml syringe was used containing 1 ml heparin to prevent clotting of the pleural fluid. The fluid was placed in EDTA treated tubes. Pleural fluid is tested for: Total Cell count, Differential cell count, Glucose, Total proteins, Lactate dehydrogenase, and Adenosine deaminase. A p-value < 0.05 was considered statistically

significant. IBM SPSS Statistics 21 manufactured by IBM USA was used for entire calculations.


**Results**

The total number of one hundred ten cases were recruited in this study in which tubercular pleural effusion cases were 64.54% and non-tubercular cases were 35.45%. Among the non-tubercular cases, 24 cases were due to malignancy, 11 were due to pneumonia, 3 cases were due to congestive cardiac failure and 1 due to rheumatoid arthritis. (Table 1, Figure 1)

Description of study of non-tuberculosis pleural effusion group and tuberculosis pleural effusion group were showed in Table 2 and 3. In the present study the mean of ADA level was 173.3, LDH level was 184.91, T. Protein was 4.80, Glucose was 112.34 and the total cell count was 3996 in tuberculosis pleural effusion group. These variables are statistically significant (<0.05). this study has clearly shown that ADA levels are significantly high in patients with tubercular pleural effusion (173.3±34.93 U/L) compared to that (20.46±7.34 U/L) in non tuberculous group.

**Table 1: Distribution of study population among pleuraleffusion cases**

Diagnosis	Number of cases (%)
<b>Tuberculous pleural effusion</b>	71 (64.54%)
<b>Non Tuberculous pleural effusion</b>	39 (35.45%)
<b>Malignancy</b>	24 (61.53%)
<b>Neumonia</b>	11 (28.20%)
<b>Rheumatoid arthritis</b>	1 (2.56%)
<b>Congestive cardiac failure</b>	3 (7.69%)



**Figure 1: Distribution of study population among pleuraleffusion cases**

**Table 2: Description of non-tuberculosis pleural effusion group**

Variables	Mean	SEM	P value
ADA	20.46±7.34	±1.84	< 0.05
LDH	81.91±6356	±7.39	< 0.05
T. protien	3.97±0.749	±.041	< 0.05
Glucose	79.47±43.26	±9.02	< 0.05
Cell Count	4872.96±1712	-	NS

**Table 3: Description of tuberculosis pleural effusion group**

Variables	Mean	SEM	P value
ADA	173.3±34.93	±3.298	< 0.05
LDH	184.91±186.3	±28.94	< 0.05
T. protein	4.80±0.642	±0.1	< 0.05
Glucose	112.34±18.98	±4.026	< 0.05
Cell Count	3996±1324.67	-	<0.05

### Discussion

This study has clearly shown that ADA levels are significantly high in patients with tubercular pleural effusion (173.3±34.93 U/L) compared to that (20.46±7.34 U/L) in non tuberculous group. Tubercular pleural effusion frequently represents a diagnostic problem even after extensive research.

Description of our study of non-tuberculosis pleural effusion group and tuberculosis pleural effusion group were showed in Table 2 and 3. In the present study the mean of ADA level was 173.3, LDH level was 184.91, T. Protein was 4.80, Glucose was 112.34 and the total cell count was 3996 in tuberculosis pleural effusion group.

These findings were similar to those observed in other studies.<sup>23</sup> However this finding is in contrast with some other studies<sup>24</sup> where they have found malignancy as the commonest cause of pleural effusion. Only a few cases of pleural effusion from congestive cardiac failure, rheumatoid arthritis and

nephrotic syndrome were noted in the present study.

Similar observations of rarity of pleural effusion by these diseases were also reported in Bangladesh.<sup>25</sup>

Other studies also showed statistically significant elevation of dehydrogenase levels in tuberculous pleural effusion compared with non-tuberculous pleural effusion.<sup>13,26-29</sup> As per the results obtained the study was discussed under the following categories. Increased adenosine deaminase (ADA) enzyme levels in tuberculous pleural effusion than non tuberculous pleural effusion. Increased lactate dehydrogenase enzyme levels in tuberculous pleural effusion than non tuberculous pleural effusion.<sup>20,30,31</sup> There is not much of a difference in total protein levels of pleural fluid in tuberculous pleural effusion and non tuberculous pleural effusion. There is not much of a difference in glucose levels of pleural fluid of tuberculous pleural effusion and non tuberculous pleural effusion. There is not much of a difference in total count of pleural fluid - but in tuberculous pleural

effusion there is lymphocytosis. ADA activity values may be increased due to other clinical entities. Valdes et al.<sup>30</sup> have also reported high levels of ADA in patients with other causes of pleural effusions (mainly lymphomas, adenocarcinomas, systemic lupus erythematosus, and pneumonia).

### Conclusion

This study has clearly shown that ADA levels are significantly high in patients with tubercular pleural

effusion compared to that in non tuberculous group. The result indicated that the analysis of ADA levels in pleural effusion constitute a very useful marker for the diagnosis of TPE which, in addition, can be made quickly in a non- invasive way. So we concluded that this is very helpful test for the diagnosis of Tubercular pleural effusion.

### References

1. Light RW: Pleural Diseases. 5th edition. Philadelphia: Lippincott Williams & Wilkins; 2007.
2. Gopi A, Madhavan SM, Sharma SK, Sahn SA: Diagnosis and treatment of tuberculous pleural effusion in 2006. *Chest* 2007, 131:880–889.
3. Porcel JM (2009) Tuberculous pleural effusion. *Lung* 187: 263–270.
4. Udhwadia ZF, Sen T (2010) Pleural tuberculosis: an update. *Curr Opin Pulm Med* 16: 399–406.
5. Baumann MH, Nolan R, Petrini M, Gary Lee YC, Light RW, et al (2007) Pleural tuberculosis in the United States: incidence and drug resistance. *Chest* 131: 1125–1132.
6. Light RW (2001) Pleural diseases. Baltimore: Lippincot, Williams and Wilkins. pp: 182–195.
7. Krenke R, Korczynski P (2010) Use of pleural fluid levels of adenosine deaminase and interferon gamma in the diagnosis of tuberculous pleuritis. *Curr Opin Pulm Med* 16: 367–375.
8. Singh R, Singh RK, Tripathi AK, Gupta N, Kumar A, Singh AK, et al. Circadian periodicity of plasma lipid peroxides and anti-oxidant enzymes in pulmonary tuberculosis. *Indian J Clin Biochem* 2004 Jan;19(1):14-20.
9. Harries AD. Tuberculosis and human immunodeficiency virus infection in developing countries. *Lancet* 1990 Feb;335(8686):387-390.
10. Raviglione MC, O'Brien RJ. Tuberculosis. In: Fauci, Braunwald, Kasper, Hauser, Longo, Jameson, Loscalzo. Eds. *Harrisons Principles of Internal Medicine*, Tata McGraw Hill, 17th ed. vol 1 p1010
11. Udhwadia ZF, Sen T. Pleural tuberculosis: an update. *Curr Opin Pulm Med* 2010 Jul;16(4):399-406.
12. Bañales JL, et al (1991). Adenosine Deaminase in the diagnosis of Tuberculous Pleural Effusion. *Chest* 99:355-57
13. Ungerer JPI, et al (1994). Significance of Adenosine Deaminase Activity and its Isoenzymes in Tuberculous effusions. *Chest* 106:33-37.
14. Seibert AF, Haynes J Jr, Middleton R, Bass JB Jr, et al (1991). Tuberculous Pleural Effusion. *Chest* 99:883-86
15. Liang QL, Shi HZ, Wang K, Qin SM, Qin XJ (2008). Diagnostic accuracy of adenosine deaminase in tuberculous pleurisy: A meta-analysis. *Resp Med* 102, 744–754.

16. Canbolat O, Ulusdoyuran S, Ozgun G, Ceyhan I, Gumuslu F, Akbay A. The comparison of adenosine deaminase activity value with polymerase chain reaction results in patients with tuberculosis. *Journal of Clinical Laboratory Analysis* 1999; 13: 209-212.
17. Riquelme A, Calvo M, Salech F, Valderama S, Pattiollo A, Arellano M, Arrese M, Soja A, Viviani P, Letelier LM, Value of adenosine deaminase (ADA) in ascitic fluid for the diagnosis of tuberculous peritonitis. *J Clin Gastroenterol* 2006; 8: 705-710.
18. Carstens ME, Burgess LJ, Maritz FJ, Taljaard JJF. Isoenzymes of adenosine deaminase in pleural effusion: a diagnostic tool? *Int J Tuberc Lung Dis* 1998; 10: 831-835.
19. Antony VB. Adenosine deaminase isoenzymes and pleural Tuberculosis. *J Lab Clin Med.*, 1996; 127: 326-327.
20. Burgess LJ, Maritz FJ, Le Roux, et al. Combined use of pleural adenosine deaminase with lymphocyte/neutrophil ratio: increased specificity for the diagnosis of tuberculous pleuritis. *Chest*, 1996; 109: 414-419.
21. Valdes L, San Jose E, Alvarez D, et al. Adenosine deaminase (ADA) isoenzyme analysis in pleural effusions; diagnostic role and relevance to the origin or increased Adenosine Deaminase in tuberculous pleurisy. *Eur Respir J*, 1996; 9: 747-751.
22. Bothamley GH. Tuberculous pleurisy and adenosine deaminase. *Thorax*, 1995; 50: 593-594.
23. Lima DM, Colares JKB and de Fonseca BAL. Combined use of the polymerase chain reaction and detection of adenosine deaminase activity on pleural fluid improves the rate of diagnosis of pleural tuberculosis. *Chest* 2003; 124: 909-914.
24. Reechaipichitkul W, Kawamatawong T, Teerajetgul Y, Patjanasoon B. Diagnostic role of pleural fluid adenosine deaminase in tuberculous pleural effusion. *Southeast Asian J Trop Med Public health* 2001; 32: 383-389.
25. Berger HW and Mejia E. Tuberculous pleurisy. *Chest* 1973; 63: 88-93.
26. Miserochi G, Agastoni E. Contents of the pleural space. *J appl Physiol.*, 1971; 30: 208-213.
27. Leuallen EC, Carr DT. Pleural effusion, statistical study of 436 patients. *N Eng J Med.*, 1955; 252: 79-83.
28. Paddock FK. The diagnostic significance of serous fluids in disease. *N ENGL J Med.*, 1940; 223: 1010-1015.
29. Lee YCG, Light RW. Adenosine deaminase for lymphocytic pleural effusions. *International pleural Newsletter*, 2004; 25-6.
30. Valdes L, Alvarez D, San Jose E, et al. Value of adenosine deaminase in the diagnosis of tuberculous pleural effusion in young patients in a region of high prevalence of Tuberculosis. *Thorax*, 1996; 50: 600-603.
31. De Oliveira HG, Rossatto ER, Prolla JC. Pleural fluid adenosine deaminase and lymphocyte proportion: clinical usefulness in the diagnosis of Tuberculosis. *Cutopathology*, 1994; 5: 27-32.