

Original article:

A clinico- bacteriological profile, antimicrobial susceptibility and outcome of neonatal sepsis in tertiary care hospital, Jaipur

***Dr. Mayur Kumar Goyal, Dr. Rohit Jain, Dr. Jaikrishan Mittal, Dr. Yogesh Vijay, Dr. Neelabh Mehru**

Department of Neonatology, Neoclinic Hospital, Jaipur, Rajasthan, India.

Corresponding author*

Abstract-

Background: Neonatal septicaemia is a clinical entity that is characterized by systemic signs and symptoms of infection and accompanied by bacteraemia in first 4 weeks of life and is one of the four leading causes of neonatal mortality and morbidity in India.

Aim: To determine the bacterial spectrum and antimicrobial susceptibility pattern of neonatal septicaemia in a tertiary care hospital of Jaipur, Rajasthan.

Materials and Methods: In this prospective observational study, 346 blood samples were collected over period of 6 months and processed from clinically suspected neonates according to standard laboratory protocol. Antimicrobial susceptibility of the isolates was done by Kirby Bauer disc diffusion method according to Clinical and laboratory standard institution (CLSI) recommendations.

Results: Blood culture reports were positive in 53 (15.31%) cases. Commonest clinical presentation of neonates with septicemia was respiratory distress and Commonest maternal risk factor was PROM >18 hours. Early onset sepsis (EOS) was present in 22.45% and late onset sepsis (LOS) in 77.55% of cases. Gram negative septicaemia (55.10%) was encountered more than Gram positive (44.90%). Coagulase negative Staphylococci (38.78%) was the predominant isolate followed by, Klebsiella spp in 34.69% cases. Best overall sensitivity among Gram-negative isolates was to polymyxin B, colistin and meropenem (100%). Gram-positive isolates had highest sensitivity to Linezolid (100 %) and Vancomycin (100%).

Conclusion: Gram negative organisms are the leading cause of neonatal septicaemia with Klebsiella spp being commonest. Coagulase negative Staphylococci is the predominant isolate among Gram positive organisms. Most of the isolates are resistant to common antibiotics.

Keywords: Neonatal sepsis, Antibiotic susceptibility, Clinical profile, Bacteriological profile

Introduction

Neonatal septicemia is a clinical entity that is characterized by systemic signs and symptoms of infection and accompanied by bacteraemia in first 4 weeks of life and is one of the four leading causes of neonatal mortality and morbidity in India¹. Incidence of neonatal septicemia in developed countries varies from 1-10/1000 live birth, whereas it is 3 times more

common in India². The incidence of neonatal sepsis according to the data from National Neonatal Perinatal Database (NNPD, 2002-03) is 30 per 1000 live births in India³. Incidence differs among hospitals depending on various factors such as obstetric and nursery practices, sex, gestational age, birth weight, out born or inborn status of babies, perinatal care, and health and nutrition of mother.⁴

Early onset sepsis (EOS) presents within the first 72 hours of life and the source of infection is generally the maternal genital tract. Knowledge about these potential risk factors would help in early diagnosis of sepsis.⁵

Late onset sepsis (LOS) usually presents after 72 hours of age. The source of infection in LOS is either nosocomial (hospital-acquired) or community-acquired.^{6,7}

Neonatal sepsis is caused by a variety of Gram-positive as well as Gram-negative bacteria, and sometimes yeasts. According to data from National Neonatal Perinatal Database (NNPD, 2002-03) in India, among intramural births, *Klebsiella pneumoniae* is the most frequently isolated pathogen (32.5%), followed by *Staphylococcus aureus* (13.6%). Among extramural neonates (referred from community/other hospitals), *Klebsiella pneumoniae* is again the commonest organism (27%), followed by *Staphylococcus aureus* (15%) and *Pseudomonas* (13%).³

Uncontrolled use of various potent and broad-spectrum antibiotics has led to emergence of resistant strains which has become a major problem in various Intensive care units.⁸ Early diagnosis and to treat neonatal infections by empirical use of antimicrobial drugs as soon as possible is must to reduce the mortality. Various diagnostic tests (hematological, biochemical and radiological) can be performed easily and results may be available in an hour or so. However, blood culture remains the gold standard for the diagnosis of neonatal septicemia.⁹

The uncertainty surrounding the clinical approach to treatment of neonatal septicemia can be minimized by periodic surveys of etiological agents and their antibiotic susceptibility patterns leading to recognition

of the most frequently encountered pathogens in a particular neonatal setting. The rational and correct use of antibiotics requires understanding of common pathogens and their drug sensitivity pattern in the regions. Due to constantly evolving antimicrobial resistant patterns there is the need for constant antimicrobial sensitivity surveillance. This will help clinicians provide safe and effective empirical therapies, develop rational prescription programs and make policy decisions and finally assess the effectiveness of all.¹⁰

Present study is conducted to determine the clinical profile, risk factors and organisms associated with bacteraemia and to find out susceptibility pattern of pathogens causing neonatal sepsis so as to provide antibiogram to NICU of Neoclinic for better patient management.

Method and material

The present prospective observational study was carried out on 346 blood samples from neonates with clinical suspicion of septicaemia, admitted in NICU of Neo clinic, Jaipur from March 2017 to September 2017.

Inclusion Criteria:

All neonates who require admission to neonatal intensive care unit (NICU) for any indication during the study period

Exclusion Criteria:

The caretaker unwilling to provide written informed consent.

The neonatal history including sex, gestational age, birth weight, term or preterm were taken. The data regarding maternal risk factor for neonatal sepsis including duration of labour, mode of delivery, maternal fever, chorioamnionitis (foul smelling liquor), maternal urinary tract infection, and duration

of rupture of membrane were collected. Data were collected in a structured proforma and were classified, analysed & evaluated by using SPSS version 21 for Windows as per aims & objectives. About 1-2 ml of venous blood was drawn by a percutaneous venous puncture following strict aseptic precautions (3 swab technique) and aseptically inoculated into blood culture bottles containing 5 -10 ml of brain heart infusion broth. Blood culture bottles were incubated at 37°C overnight aerobically. Primary subcultures were done after 24 hours of incubation on to Blood agar & Mac- Conkey agar. If no growth occurred on plates after overnight incubation, bottles were incubated further & followed up by examining the broth daily and doing a final subculture at the end of day 7 or at appearance of signs of growth, whichever was earlier. The positive growth was identified by conventional methods according to the standard laboratory protocol, including colony morphology, Gram's staining & biochemical reactions. After the identification of bacteria, antimicrobial susceptibility testing was done by Kirby- Bauer disc diffusion method on Muller Hinton agar as per Clinical and Laboratory Standards Institute (CLSI) guidelines.¹¹ The antibiotics which were used in our study were based on the standard protocol of the hospital and departmental policies (as per CLSI).

The clinical signs suggestive of infection include:⁵

- Refusal to feed, poor cry, lethargy
- Temperature instability [Fever (>37.5°C), hypothermia (< 36.5 °C)]
- Off color appearance, dusky peripheries
- Prolonged CFT (>3sec)
- Apnea, worsening respiratory distress
- Feed intolerance, abdominal distension
- Blood in stool and other bleeding manifestation

The perinatal risk factors include:⁵

- Low birth weight (<2500 grams) or prematurity
- Febrile illness in mother with evidence of bacterial infection within in 2 weeks prior of delivery
- Rupture of membrane >18 hours
- Prolonged labor (sum of 1st and 2nd stage of labor ≥24 hours)
- Single unclean or >3 sterile vaginal examinations during labor
- Foul smelling liquor
- Perinatal asphyxia (Apgar score <4 at 1 min)

Presence of foul smelling liquor or three of the above mentioned risk factors warrant initiation of antibiotic treatment. Infants with two risk factors were investigated and then treated accordingly.

Various predisposing factors to increase risk of nosocomial sepsis include:

- Low birth weight
- Prematurity
- Admission in intensive care unit
- Mechanical ventilation
- Invasive procedures
- Administration of Parenteral fluids, and
- Use of stock solutions

Results:

During the study period, a total of 346 neonates with clinical sepsis were studied. The various baseline characteristics of the study including risk factor for neonatal sepsis are included Blood culture positivity was found in 53 cases (15.32%) while 293 (84.68%) were blood culture negative. Bacterial species were found in 49 (92.45%) and Candida species in 4 (7.55%) cases.

Out of 53 culture positive cases, Incidence was 38 (16.24%) male and 11 (9.82%) female neonates with the male-to-female ratio of 3.45:1. Late onset sepsis

(LOS) cases were found to be higher than Early onset sepsis (EOS). 11 neonates (22.45%) had EOS and 38 (77.55%) had LOS. Incidence of septicaemia was higher in preterm neonates (18.84%) as compared to term (11.06%).

Among culture positive cases, 19 (15.57%) were of low birth weight, 6 (19.35%) were of very low birth weight, and 4 (25%) were of extreme low birth weight. Normal birth weight was found in 20 (11.29%) cases. So incidence of neonatal septicaemia was higher in very low birth weight neonates compared to normal birth weight.

Of the various maternal risk factors affecting neonatal septicaemia, rupture of membranes for more than 18 hours was the commonest factor with 63.64% cases followed by foul smelling liquor in 27.27%, febrile illness in mother 18.18% and maternal urinary tract infection (UTI) in 9.09%.

According to clinical presentation, respiratory distress (65.30%) was commonest presentation among culture positive cases followed by encephalopathy (22.44%) and seizures (14.28%).

Out of 49 cases, 83.67 % baby survived and discharged while 10.20% expired and 6.13% baby was LAMA.

Etiology of the 49 bacterial culture positive isolates included Gram-negative bacilli (27/49, 55.10%) and Gram positive cocci (22/49, 44.90%). Coagulase Negative Staphylococci (38.78%) was the commonest bacterial isolate followed by Klebsiella spp (34.69%), Acinetobacter (14.29%), Enterobacter (4.08%), Enterococcus spp (4.08%), Coagulase positive Staphylococci (2.04%) and Escherichia coli(2.04%).

Gram positive organisms showed highest sensitivity to Linezolid (100 %) and Vancomycin (100 %), followed by Doxycycline (86.36%), Meropenam (86.36 %) and Gentamycin (81.82 %) and least sensitivity to Cefuroxime (4.54 %).

Gram negative organisms showed good sensitivity to polymyxin B, colistin and meropenem i.e 100%, 96.27% and 88.88% respectively. Ceftazidime showed poor sensitivity amongst Gram negative organisms.

Table- 1: Incidence of Neonatal septicemia

Cases	Number of Cases (n= 346)	Percentage (%)
No Sepsis	180	52.02 %
Suspected Sepsis	113	32.66%
Culture proven sepsis	53	15.32%

Table-2: Gender distribution of bacterial culture positive cases

Sex	Total Cases (N=346)	Number of culture positive cases (n=49)	Percentage (%)
Male	234	38	16.24%
Female	112	11	9.82%

Table- 3: Distribution of incidence of bacterial culture positive cases according to Birth Weight

Birth weight	Total Cases (N=346)	Number of culture positive cases (n=49)	Percentage (%)
>2.5 kg	177	20	11.29%
1.5-2.5 kg	122	19	15.57%
1-1.5 kg	31	6	19.35%
<1 kg	16	4	25%

Table-4: Distribution of incidence of gestational age in bacterial culture positive cases

Gestational age	Total Cases (N=346)	Number of culture positive cases (n=49)	Percentage (%)
Full term (≥ 37 weeks)	208	23	11.06%
Pre term (<37 weeks)	138	26	18.84%

Table-5: Distribution of bacterial culture positivity in Early & Late onset cases

Age at onset of symptoms	Number of cases(n=49)	Percentage (%)
Early onset septicemia	11	22.45%
Late onset septicemia	38	77.55%

Table-6: Distribution of maternal risk factors (Early Onset Sepsis n=11) in bacterial culture positive cases

Maternal Parameters	Cases (n=11)	Percentage (%)
Prolonged rupture of membrane (PROM)> 18 Hrs	7	63.64 %
Maternal Urinary tract infection	1	9.09 %
Febrile illness in mother	2	18.18 %
Foul Smelling Liquor/ features of chorioamnionitis	3	27.27 %

Table- 7: Distribution of bacterial culture positive cases in relation to clinical presentation of Neonates

Clinical presentation	Number of cases(n= 49)	Percentage (%)
Respiratory distress	32	65.30 %
Seizures	7	14.28 %
PA / Encephalopathy	11	22.44 %
Dullness	6	12.24 %
Poor feeding	5	10.20 %
Fever	4	8.16 %
Excessive cry	2	4.09 %
Bleeding	3	6.12 %

Table-8: Spectrum of bacterial isolates

Isolates	Number of cases (n=49)	Percentage (%)
CONS	19	38.78%
Klebsiella	17	34.69%
Acinetobacter	7	14.29%
Enterobacter	2	4.08%
Enterococcus	2	4.08%
COPS	1	2.04%
E coli	1	2.04%

Table- 9: Antibiotic sensitivity pattern of Gram positive bacteria (n=22)

Antibiotic	Number of cases	Percentage (%)
Amikacin	15	68.18%
Amoxicillin/ Clavulanic acid	2	9.09%
Ampicillin	2	9.09%
Ampicillin / Sulbactam	4	18.18%
Azithromycin	5	22.73%
Cefoxitin	12	54.54%
Ciprofloxacin	7	31.82%
Cefuroxime	1	4.54%
Gentamycin	18	81.82%
Levofloxacin	7	31.82%
Linezolid	22	100%
Ofloxacin	8	36.36%
Meropenam	19	86.36%
Teicoplanin	17	77.27%
Vancomycin	22	100%
Doxycyclin	19	86.36%

Table- 10: Antibiotic sensitivity pattern of Gram negative bacteria (N=27)

Antibiotic	Number of cases(n)	Percentage (n?N%)
Amikacin	14	51.85%
Colistin	26	96.27%
Ceftazidime + clavulamic acid	2	7.41%
Ceftriaxone	5	18.52%
Imipenam	22	81.48%
Meropenam	24	88.88%
Ciprofloxacin	13	48.15%
Polymyxin B	27	100%
Gentamycin	7	25.92%
Levofloxacin	14	51.85%
Piperacillin / Tazobactam	11	40.74%
Piperacillin	5	18.52%
Cefipime	7	25.92%
Tobramycin	10	37.04%

Table-11: Outcome of blood culture proven bacterial sepsis cases (n=49)

Outcome	Number of cases	Percentage (%)
Discharge	41	83.67%
LAMA	3	6.13%
Expired	5	10.20%

Discussion:

The present study was carried out in the Department of neonatology, Neo clinic, Jaipur. Total 346 babies admitted during study period were enrolled and out of which 180 (52.02%) cases had no sepsis, 113 (32.65%) had suspected sepsis, 22 (6.35 %) had Gram positive sepsis, 27 (7.80%) had gram negative sepsis and 4 (1.11 %) cases had fungal sepsis. Out of 346 neonates, 53 were found to be blood culture positive, giving a culture positivity rate of 15.31%.

Culture positivity

According to previous studies blood culture positivity in neonatal septicaemia varies from 18.8%⁴⁸ to 64.87%¹³. Our results are comparable with studies conducted by P.Jyothi et al¹⁴, Sanjay Rathod et al¹⁵, and Ramesh bhat et al¹⁶ Low positivity might be due to administration of antibiotic before blood collection, possibility of infection with anaerobes or presence of fastidious organisms and quality controlled care in our NICU.

However, positivity rate in our study is low as compared to the results reported by Roy et al¹⁷, Hirah shah et al¹⁸ and Bheemasamudra et al¹⁹, which, might be attributed to the better infection control practices in our Neonatal ICUs.

Demography

Sex

In the present study males(16.24%) were affected more than females(9.82%) with male to female ratio of 1.65:1. Similar findings were also observed by Kavitanimbooret al.⁴, Anitha Sharma et al²⁰. Deepandra garg et al²¹ and Bheemasamudra et al¹⁹. Male preponderance may be linked to the X linked

immunoregulatory gene resulting in susceptibility to infections.

Preterm Vs Term-

Prematurity is an important neonatal factor predisposing to infection. Preterm neonates are more prone to septicaemia because of the immaturity of immune system as compared to full term babies. In our study, blood culture positivity was found more in preterm neonates (18.84%) than full term (11.06%). Our results are similar to other studies like Bheemasamudra et al¹⁹ and Arpita Jigar Shah et al²². However, surprisingly in a study by Dechen C Tsering et al²³ positivity was higher in full term as compared to preterm.

Birth weight

Khatua et al²⁴ and Mondal et al²⁵ stated that low birth weight neonates have low IgG level and are more susceptible to infections. According to Barbara J. Stoll²⁶, rate of infection is inversely proportional to birth weight. In our study septicaemia was more common in extreme low birth weight neonates (<1000 gm) (25%) while babies weighing >2.5 Kg comprising of 11.29% cases. Our findings were consistent with Bheemasamudra et al.¹⁹

Early Vs Late Onset Septicemia

Although incidence of EOS is more than LOS due to immature immunologic response of neonates in the first week of life, making them more susceptible to infection in this period. We found in our study that LOS(77.55 %) was more common than EOS(22.45 %), Our findings are also compatible with the reports from the other developing countries like higher incidence of LOS reported by Daynia

et al²⁷ and Kurien Kuruvilla²⁸. On contrary, EOS was more common in studies done by Kaushal et al²⁹, Sweetha nayak et al³⁰ and P. Jyothi et al¹⁴.

EOS cases might be less in number due to administration of intrapartum antibiotics prophylaxis and Institutional deliveries. Most of the babies having late onset septicemia admitted on later days of life and might infected from surrounding before admitted in NICU.

Clinical Presentation

Of the various clinical presentations of neonates with septicaemia, our study revealed that respiratory distress was the commonest presentation followed by dullness. Study by Kavita nimboor⁴ and Deepandra garg et al²¹ showed refusal to feed being the commonest symptom followed by respiratory distress. Mamta jagoo et al³¹ reported refusal to feed as leading symptom followed by respiratory distress. Khatua et al²⁴ reported refusal to feed, hypothermia and respiratory distress as common clinical presentation.

Spectrum of organisms

Gram-negative organisms are more common pathogens and are mainly represented by Klebsiella, Escherichia coli, Pseudomonas, Enterobacter spp and Acinetobacter spp. Among the Gram-positive organisms, Staphylococcus aureus, CoNS, Streptococcus pneumoniae, and Streptococcus pyogenes are common isolates.¹⁴

Gram negative septicaemia (55.10 %) was encountered more than Gram positive (44.90 %) in the present study which is comparable to studies conducted by Tsering DC et al²³, Arpita Jigar et al²² and P. Jyothi et al¹⁴. The increased susceptibility of neonates to the Gram negative bacteria may be explained by the fact that the antibodies against these organisms are primarily IgM type, which do not transfer passively through placenta and are at very low level in blood (about 5% of adult value) at birth, and reaches the adult level by 2 years of age. This is in contrast

with IgG type, which are passively transferred to placenta and are almost at adult level at birth and falls gradually reaching lowest level around 3 to 4 months of age after which they start to rise again gradually. Adequate IgG (except IgG 2-subtype) levels at term, afford protection against several Gram positive bacteria.³²

In the present study Coagulase negative Staphylococci (38.78%) was the predominant isolate followed by Klebsiella spp in 34.69 % cases. CoNS is normally considered as a skin contaminant when isolated from blood. The presence of this bacteria in blood in critically ill babies should be considered as significant and should be treated. In our study, we considered CoNS as a true pathogen because the antibiotic sensitivity pattern of our CoPS and CoNS isolates was almost similar and they correlated clinically. Moreover the samples were collected with proper sterile precautions, so the chances of contamination with skin flora were minimal.

Among Gram negative bacteria, we found Klebsiella spp (62.96%) as commonest followed by Acinetobacter (25.93%), similar to study by K.Chug et al³³, Anuradha de et al¹², and Anitha sharma et al²⁰. This was in concordance with National Neonatal Perinatal Database (NNPD- 2003)³ with Klebsiella spp as a predominant Gram negative isolate associated with neonatal sepsis. However in some studies by S.G Joshi et al³⁴, Tsering DC et al²³, and Ramesh bhat et al¹⁶, Pseudomonas spp was the commonest isolate among Gram negative bacteria.

In our study Coagulase negative Staphylococci was the commonest (86.36%) isolate among Gram positive cocci followed by Enterococcus spp (9.09%) and Coagulase positive Staphylococci (4.55%). This is in accordance to study by Edwin dias et al³⁵, Daynia E et al³⁶, G.P Mondal et al³⁷ and Isaacs D et al³⁸.

Antimicrobial susceptibility

The antimicrobial susceptibility (AST) pattern differs in different studies as well as at different times in the same hospital in Indian and overseas studies because of the different hospital based antibiotic policies.

In our study AST revealed that majority of Gram negative bacteria were sensitive to polymyxin B (100%), colistin (96.27%), meropenem (88.88%) and imipenem (81.48%). P Jyothi et al¹⁴ also found best overall sensitivity among Gram negative isolates to imipenem (93%). Among aminoglycosides maximum sensitivity was seen to amikacin (57.78%) which is in line with Madavi et al³⁹, who reported 60.60% sensitivity to amikacin. Least sensitivity was seen to extended spectrum cephalosporins in our study. Among Gram negative bacteria, the predominant *Klebsiella* spp showed good sensitivity to polymyxin B (100%), colistin (100%), meropenem (94.12%) and imipenem (88.23%), which is similar to results reported by Kaushal seth et al²⁹ and Maimoona mustafa et al⁴⁰ with 100% sensitivity to imipenem. Sensitivity to amikacin (52.94%) was better than gentamycin (35.29%) among amino-glycosides in our study which was comparable to results of studies by Bheemasamudraet al¹⁹, Ramesh bhat et al⁴¹ and Maimoonamustafa et al⁴⁰. In our study Gram positive

bacteria showed good sensitivity to Linezolid (100%), vancomycin (100%), meropenem (86.36%), doxycycline (86.36%) and Gentamycin (81.82%) which is comparable to study by Mamtajagoo et al³¹, Arpita shah et al²², and Kaushal v et al²⁹. Sensitivity to ampicillin (9.09%), amoxicillin and clavulanic acid (9.09%) and cefuroxime (4.54%) was lowest among Gram positive bacteria in our study.

Carbapenems (imipenem and meropenem), colistin and polymyxin B were found to be the most potent antimicrobial agents against Gram negative organisms. Majority of Gram positive isolates were sensitive to linezolid and vancomycin.

Outcome of neonatal septicemia-

Out of 49 babies, 23 (46.94%) babies discharged, 18 (36.73%) went Discharge on request, 3 (6.13%) left against medical advice and 5 (10.20%) babies expired as a consequence of the underlying disease. Iyeret al⁴² also showed similar outcome in his study.

Conclusion

Gram negative organisms are the leading cause of neonatal septicaemia with *Klebsiella* spp being commonest. Coagulase negative Staphylococci is the predominant isolate among Gram positive organisms. Most of the isolates are resistant to common antibiotics.


BIBLIOGRAPHY

1. Shankar MJ, Aggarwal R, Deorari AK, Paul VK. Symposium on AIIMS protocol in Neonatology III. Indian J Paediatr, 2008;75:261-6.
2. Singh M. Care of the newborn; 6th edition; Meharban Singh, Sagar Publications; page 212 220.)
3. Report of the National Neonatal Perinatal Database (National Neonatology Forum) 2002-03.
4. Kavita nimboor. "Bacteriological (aerobic) study of neonatal septicemia in NICU."
5. H.Takkar VP, Bhakoo ON, Narang A. Scoring system for the prediction of early neonatal infections. Indian Pediatr, 1974;11:597-600
6. Baltimore RS. Neonatal nosocomial infections. Semin Perinatol 1998;22: 25-32.
7. J. Wolach B. Neonatal sepsis: pathogenesis and supportive therapy. Semin Perinatol, 1997;21:28-38


8. Sanjay D Rathod, Palak V Bhatia, Parimal H Patel, Jayshri D Pethani1, Lata R Patel, Bimal Chauhan. National Journal Of Medical Research, 2012; Print Issn: 2249 4995 | Eissn: 2277 8810 Volume 2 | Issue 4 | Page 467 .
9. Sharma A, Kuty CV, Sabharwal U, Rathee S, Mohan H. Evaluation of sepsis screen for diagnosis of neonatal septicemia. Indian J Pediatr, 1993; 60: 559-563
10. Karki S1, Rai GK2, Manandhar R3. Bacteriological Analysis and Antibiotic Sensitivity Pattern of Blood Culture Isolates in Kanti Children Hospital. J. Nepal Paediatr. Soc. 2010; Vol 30 ;94-97
11. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing; 23rd informational supplement. Pennsylvania: Clinical and Laboratory Standards Institute; 2013.
12. De Anuradha et al., "Bacteraemia in Hospitalized children – A one Year Prospective study". IJMM: 1995; 13: 72-75.
13. Tallur SS, KasturiAV, NadigrSD, KrishnaBV. Clinico - bacteriological study of neonatal septicemia in Hubli. Indian J Pediatr, 2000;67:169-74.
14. Jyothi, et al.: Bacteriological profile of neonatal septicemia Journal of Natural Science, Biology and Medicine, July 2013; Vol 4 , Issue 2, IP: 106.219.29.67
15. Philp A G S, Neonatal meningitis in the New Millennium. American Academy of Pediatrics, 2003; 4(3) : e73.
16. Ramesh Bhat Y, Leslie Edward S Lewis and Vandana KE. Bacterial isolates of early-onset neonatal sepsis and their antibiotic susceptibility pattern between 1998 and 2004: an audit from a center in India. Italian Journal of Pediatrics ,2011, 37:32
17. Roy I, Jain A, Kumar M, Agarwal S K. Bacteriology of neonatal septicaemia in a tertiary care hospital of northern India. Indian J Med Microbiol, 2002;20:156-9.
18. Dr. Hiral Y Shah et al. Antibiotics Susceptibility Patterns of Bacterial Isolates Among Neonatal Septicemia in Tertiary Care Hospital, Jamnagar, Gujarat. Global Research Analysis, 2012; 1 (5) : 2277 - 8160
19. Rajendraprasad BM, Basavaraj KN, Antony B. Bacterial spectrum of neonatal septicemia with their antibiogram with reference to various predisposing factors in a tertiary care hospital in Southern India. Ann Trop Med Public Health, 2013;6:96-9.
20. Sharma Anitha et al., "Diagnostic and prognostic role of CRP and mESR in neonatal septicaemia". Indian paediatrics, 1993; 30: 347-349.
21. Dr. Deepandra Garg, Dr. Neha Agrawal. Aetiology And Presentation Of Neonatal Septicaemia At Tertiary Care Hospital Of Southern Rajasthan. International Journal of Medical Science and Education, Jan-March 2014 ; Vol.1; Issue: 1.
22. Arpita Jigar Shah, Summaiya A. Mulla, and Sangita B. Revdiwala. JClinNeonatology. 2012 Apr-Jun; 1(2): 72–75.doi: 10.4103/2249-4847.96753
23. Tsering DC, Chanchal L, Pal R, Kar S. Bacteriological profile of septicemia and the risk factors in neonates and infants in Sikkim. J Global Infect Dis, 2011;3:42-5.
24. Khatua SP, Das AK, Chatterjee BD, Khatua S, Ghose B et al. Neonatal septicaemia. Indian J Pediatr, 1986; 53: 509-14.

25. Ghotaslou R, Ghorashi Z, Nahaei MR. Klebsiella pneumoniae in neonatal sepsis: A 3-year study in pediatric hospital of Tabriz, Iran. *JPN J Infect Dis*, 2007;60:126–8.
26. Shrestha S, Adhikari N, Rai BK, Shreepaili A. Antibiotic Resistance Pattern of Bacterial Isolates in Neonatal Care Unit. *J Nepal Med Assoc*, 2010;50:277–81.
27. Daynia E. Ballot, Trusha Nana, Charlotte Sriruttan, and Peter A. Cooper Bacterial Bloodstream Infections in Neonates in a Developing Country.
28. Kurien Anil Kuruvilla, Swati Pillai, Mary Jesudason* and Atanu Kumar Jana,” Bacterial Profile of Sepsis in a Neonatal Unit in South India”, *Indian Paediatrics*, 1998; 35:851-858.
29. Kaushal V. Sheth, Tejas K. Patel, C.B. Tripathi. *Asian J Pharm Clin Res*,2012; Vol 5, Issue 3, 46-50.
30. Nayak S, Rai R, Kumar VK, Sanjeev H, Pai A, Ganesh H R. Distribution of microorganisms in neonatal sepsis and antimicrobial susceptibility patterns in a tertiary care hospital . *Arch Med Health Sci*, 2014;2:136-9.
31. Jajoo M, Kapoor K, Garg L K, Manchanda V, Mittal S K. To study the incidence and risk factors of early onset neonatal sepsis in an out born neonatal intensive care unit of India. *J ClinNeonatology* , 2015;4:91-5.
32. Singh MB. Perinatal infections. In: *Care of the newborn*. 6th ed. New Delhi: Sagar publications, 2004; p.196-216.
33. Chug. K. et al., “Bacteriological profile of neonatal septicaemia”. *Indian J. Pediatr*,1988; (55) 961-965.
34. Joshi S.G. et al., “Neonatal Gram negative bacteraemia”. *Indian J. Pediatrics*, 2000; 67: 27: 32.
35. Edwin dias and vigneshwaran p. the bacterial profile of neonatal septicaemia in a rural hospital in South India. *Journal of Clinical and Diagnostic Research*. 2010; 4:3328-3331.
36. Sohn AH, Garrett DO, Sinkowitz-Cochran RL, Grohskopf LA, Levine GL, Stover BH, et al. Prevalence of nosocomial infections in neonatal intensive care unit patients: results from the first national point prevalence survey. *J Pediatr* ,2001; 139: 821-7.
37. Mondal G.P. et al. “Neonatal Septicemia among in born and outborn babies in referral Hospital”. *Indian J. Pediatr*, 1991; 58: 529-533.
38. Isaacs D et al. Late onset infections in neonatal units” *J. Paediatr Child Health*. 1996; 32(2): 158-61.
39. Madaviet. al. Clinico-bacteriological profile and antibiotic sensitivity pattern of neonatal septicaemia- a prospective observational study. *Int J Cur Res*. 2015; 7 : 5.
40. Maimoona Mustafa and Syed LaeeqAhmed. Bacteriological profile and antibiotic susceptibility patterns in neonatal septicemia in view of emerging drug resistance. *J Med Allied Sci*, 2014; 4 (1): 02-08.
41. Ramesh Bhat Y, Leslie Edward S Lewis and Vandana KE. Bacterial isolates of early-onset neonatal sepsis and their antibiotic susceptibility pattern between 1998 and 2004: an audit from a center in India. *Italian Journal of Pediatrics*, 2011, 37:32
42. Iyer CR et al. clinical profile and outcome of neonates with suspected sepsis from a rural medical college hospital of south india. *Int j Contempediater*. 2018 Jan;5(1):55-60.


Graph-1: Incidence of Neonatal septicemia (N= 346)



Graph- 2: Spectrum of bacterial isolates (N= 49)



Graph-3: Antibiotic sensitivity pattern of Gram positive bacteria (n=22)



Graph-4: Antibiotic sensitivity pattern of Gram negative bacteria (N=27)

